

# Reversing the temporal envelope of a heralded single photon using a cavity

Bharath Srivathsan,\* Gurpreet Kaur Gulati,\* Alessandro Cerè,\* Brenda Chng,\* and Christian Kurtsiefer†

(Dated: September 4, 2014)

We demonstrate a way to prepare single photons with a temporal envelope that resembles the time reversal of photons from the spontaneous decay process. We use the photon pairs generated from a time-ordered ~~atomic~~ cascade decay: the detection of the first photon of the cascade is used as a herald for the ground-state transition resonant second photon. We show how the interaction of the heralding photon with an asymmetric Fabry-Perot cavity reverses the temporal shape of its twin photon from a decaying to a rising exponential envelope. This single photon is expected to be ideal for interacting with two level systems.

Absorption of a single photon by a single atom or an ensemble of atoms is an interesting problem from a fundamental point of view, and is also essential for many quantum information protocols [1–4]. One of the requirements for an efficient absorption is that the temporal shape of the incident photon is the time reversal of the photon from the spontaneous decay process [5, 6]. Temporally shaped light pulses have been utilized in many recent experiments to achieve efficient interactions between light and matter [7, 8]. In particular, the advantage of using a rising exponential shaped single photon for absorption in an atomic ensemble was demonstrated in [9], and shaped multiphoton pulses for exciting a single atom was demonstrated in [10]. [Interaction of single photons with other two-level systems such as quantum dots \[11, 12\], single molecules \[13\] and super-conducting circuits \[14\] are also currently being researched and require a “time reversed” envelope of the photons for efficient interaction.](#)

Efficient preparation of single photons with narrow bandwidth and a rising exponential envelope is not trivial. One solution is the direct modulation of a heralded photon generated by an atomic medium [15]. This technique results in unavoidable losses due to filtering. We have previously demonstrated a scheme to generate single photons with a rising exponential shape by heralding on photon pairs produced by cascade decay [16] without filtering. The drawback of this scheme is that the photon with the rising exponential envelope is not resonant with an atomic ground state transition.

In this letter, we combine the asymmetric cavity design used by Bader et al. [17] with the well known temporal correlation properties of photon pairs [18] to invert the temporal envelope of the generated photon pairs: with the proper heralding sequence we obtain a rising exponential single photon resonant with a ground state transition of  $^{87}\text{Rb}$ . [This technique can be extended to other physical systems with a cascade level structure to obtain these photons \[19–21\]. Similar ideas have been used in the past for non-local dispersion cancellation and ghost imaging \[22, 23\].](#)

The photons emerging from an atomic cascade decay have a well defined time order. The first photon of the cascade (signal) is generated before the photon resonant with the ground state (idler). The resulting state can be

described by a two photon wave function [24] of the form

$$\psi(t_s, t_i) = A e^{-(t_i - t_s)/2\tau} \Theta(t_i - t_s), \quad (1)$$

where  $t_s, t_i$  are the detection times of the signal and idler photons, and  $\Theta$  is the Heaviside step function. In this notation, the probability of observing a pair is proportional to  $|\psi(t_s, t_i)|^2$ . The exponential envelope and the decay time  $\tau$  is a consequence of the atomic evolution of the cascade decay. If the detection of a signal photon is used as herald, the idler mode has a single photon state with a exponentially decaying temporal envelope starting at  $t_i = t_s$ . Similarly, if the detection of an idler photon acts as a herald, the signal photon has an exponentially rising temporal envelope.

An asymmetric cavity with the appropriate parameters transforms a light field with an exponentially rising envelope into one with a decaying envelope [17]. The main point of this work is that coupling the signal mode into a properly tuned asymmetric cavity before heralding results in a “time reversed” envelope for the idler photon.

The asymmetric cavity is formed by a partially reflective mirror  $M_1$ , and a highly reflective mirror  $M_2$ , see Fig. 1. The effect of the cavity on the signal mode is described as a frequency-dependent phase factor [25, 26],

$$C(\delta') = \frac{\sqrt{R_1} - \sqrt{R_2} e^{i\delta'/\Delta\nu_f}}{1 - \sqrt{R_1 R_2} e^{i\delta'/\Delta\nu_f}}, \quad (2)$$

where  $R_{1,2}$  are the reflectivities of  $M_1$  and  $M_2$ ,  $\Delta\nu_f$  is the free spectral range of the cavity, and  $\delta'$  the detuning from the cavity resonance. For  $R_2 = 1$ , the transformation of the incoming mode is lossless, i.e.,  $|C(\delta')| = 1$ .

The cavity transforms the two-photon wavefunction in Eq. (1) into the two-photon wavefunction  $\tilde{\psi}(t_s, t_i)$ :

$$\tilde{\psi}(t_s, t_i) = \mathcal{F}_s^{-1} [C(\omega_s - \omega_s^0 - \delta) \cdot \mathcal{F}_s [\psi(t_s, t_i)]] , \quad (3)$$

where  $\mathcal{F}_s$  denotes a Fourier transform from  $t_s$  to  $\omega_s$ , and  $\delta$  is the detuning of the cavity resonance from the signal photon center frequency  $\omega_s^0/2\pi$ .

If the ring-down time of the cavity matches the coherence time  $\tau$  of the photon pair in Eq. (1), the resulting wavefunction is:

$$\tilde{\psi}(t_s, t_i) = \frac{A}{\sqrt{1 + 4\delta^2\tau^2}} [ 2\delta\tau e^{-(t_i - t_s)/2\tau} \Theta(t_i - t_s)$$




FIG. 1: Schematic of the four-wave mixing experiment in collinear geometry. IF<sub>1</sub>, IF<sub>2</sub>: interference filters, used to combine pump beams and to separate the photons pairs. SMF: Single mode optical fibers. M<sub>1</sub>, M<sub>2</sub>: cavity mirrors. The incoming and outgoing mode of the cavity are separated by a polarising beam splitter (PBS) and a quarter wave plate ( $\lambda/4$ ). D<sub>1</sub>, D<sub>2</sub>: silicon avalanche photodiodes (APD). The inset shows the cascade level scheme for generation of photon pairs in  $^{87}\text{Rb}$ .

$$+e^{(t_i-t_s)/2\tau} \Theta(-t_i+t_s)] \quad (4)$$

with an exponentially rising and an exponentially decaying component. Their relative weight can be controlled by the detuning  $\delta$ , and for  $\delta = 0$ , a time-reversed version of Eq. (1) is obtained:

$$\tilde{\psi}(t_s, t_i) = A e^{(t_i-t_s)/2\tau} \Theta(-t_i+t_s). \quad (5)$$

Heralding on the detection of the modified signal photon results in an idler photon state with a rising exponential envelope, ending at  $t_i = t_s$ . The cavity thus effects a reversal of the temporal envelope of the heralded idler photons from an exponential decay to a rise.

To experimentally investigate this method, we used the setup shown in Fig. 1. We generate time-ordered photon pairs by four-wave mixing in a cold ensemble of  $^{87}\text{Rb}$  atoms in a cascade level scheme. Pump beams at 795 nm and 762 nm excite atoms from the  $5S_{1/2}, F = 2$  ground level to the  $5D_{3/2}, F = 3$  level via a two-photon transition. The 776 nm (signal) and 780 nm (idler) photon pairs emerge from a cascade decay back to the ground level and are coupled to single mode fibers. All four modes are collinear and propagate in the same direction. The coherence time  $\tau$  of the photon pairs is determined by a time-resolved coincidence measurement between the detection of signal and idler photons to be 5.9 ns. Details about the photon pair source can be found in [27] and [16].

One of the modes of the photon pairs (signal in Fig. 1) is coupled to the fundamental transverse mode of an




FIG. 2: Transformation of the temporal envelope of the heralded idler photon from exponential decay to rise when the cavity is in the signal mode. The y-axis shows the coincidence rate  $G_{si}^{(2)}$  between the detectors D<sub>1</sub> and D<sub>2</sub> as a function of the detection time difference. The dashed lines represent  $|\tilde{\psi}(t_s, t_i)|^2$ , obtained from the model described by Eq. (4) for the indicated cavity detunings  $\delta$ , with amplitude  $A$  as the only free parameter used to fit the experimental points.

asymmetric cavity, formed by mirrors M<sub>1</sub>, M<sub>2</sub> with radii of curvature of 100 mm and 200 mm, respectively. We characterize the cavity using a frequency stabilized laser of wavelength 776 nm. The reflectivity of mirror M<sub>1</sub> is determined by direct measurement with a PIN photodiode to be  $R_1 = 0.9410 \pm 0.0008$ . Transmission through the mirror M<sub>2</sub> and absorption by the mirrors leads to losses in the cavity. The loss per round trip is determined from the transmission through and the reflection from the cavity and is included in the reflectivity of M<sub>2</sub>,  $R_2 = 0.998 \pm 0.001$ . The mirrors are separated by 5.5 cm corresponding to a free spectral range  $\Delta\nu_f = 2.7$  GHz. Therefore, an incident photon of Fourier bandwidth  $1/(2\pi\tau) = 27$  MHz interacts effectively with only one longitudinal mode of the cavity, ensuring that Eq. (2) is an adequate model. The light reflected off the cavity is separated from the incident mode by using a polarising beam splitter (PBS) and a quarter waveplate ( $\lambda/4$ ).

We infer the temporal shape of the heralded photons from the time distribution of the coincidence rate  $G_{si}^{(2)}$  between the APDs D<sub>1</sub> and D<sub>2</sub> (time resolution  $< 1$  ns). In Fig. 2 we show  $G_{si}^{(2)}$  for three different cavity-




FIG. 3: Temporal envelope of the idler photon when the cavity is aligned and tuned to resonance with the idler mode. The dashed line represents  $|\tilde{\psi}(t_s, t_i)|^2$  obtained from Eq. (3) by swapping  $i$  and  $s$ . Also in this case, the amplitude is the only free parameters used in the fit.

photon detunings. When the cavity resonance is tuned far away from the signal photon frequency  $\omega_s^0$ , in this case about  $\delta/2\pi = 120$  MHz, the temporal envelope remains nearly unchanged from the exponential decay obtained without cavity. [Off-resonant coupling of the incident signal photon to the cavity leads to the residual coincidences at times  \$t\_i - t\_s < 0\$ .](#) At  $\delta/2\pi = 27$  MHz, the time distribution becomes a symmetric exponential, and on resonance,  $\delta/2\pi = 0$ , we obtain a rising exponential shape. For the three detunings the measurement agrees with the shape expected from Eq. (4): the exponential time constants remain unchanged and the new temporal shapes are determined by the phase shift across the cavity resonance via Eq. (2).

From the time distribution of the coincidence counts it is evident that the situation is symmetrical to what we presented in [16]: by heralding on the signal photon we now obtain an idler photon with a rising exponential temporal envelope. This result, though predicted by the theory, is particularly exciting: the idler photon is resonant with a ground state transition and the obtained temporal envelope is similar to the time reversal of the one obtained by spontaneous decay. The only deviation from the predicted shape occurs for a short time interval after the detection of the herald ( $t_i - t_s > 0$ ). We attribute this deviation to an imperfect matching between the signal and cavity spatial modes.

To confirm the predictive power of our model, we repeated the same experiment swapping the roles of the signal and idler modes. This corresponds to swapping the subscripts  $s$  and  $i$  in Eqs. (3) and (4). Figure 3 shows the time resolved coincidence rate  $G_{si}^{(2)}$  between the signal and modified idler photons with the cavity tuned to resonance with the idler central frequency. In this case the




FIG. 4: Mean photon number in the cavity estimated using Eq. (6). On the left, the detection of an idler photon is used as herald and the cavity is in the signal mode. In this case we observe the interaction of an exponentially rising waveform with the cavity. On the right, the roles of signal and idler are swapped and the cavity interacts with an exponentially decaying incident photon.

cavity transforms the exponentially rising temporal envelope into a more complex shape. Our model describes accurately this complex shape, as can be seen from the dashed line in Fig. 3.

Using the same setup, we can infer the population of the cavity mode as a function of time, and observe its dependence on the envelope of the incident photon. We estimate the mean photon number  $\langle n(t) \rangle$  in the cavity using an algorithm similar to the one presented in [17]. We compare the time distributions of coincidence rates  $G_{fr}^{(2)}$  and  $G_{or}^{(2)}$  when the cavity is tuned far-off resonance ( $\delta/2\pi = 200$  MHz) and on resonance ( $\delta/2\pi = 0$  MHz) with the incident photons, normalized against the total far-off resonance coincidences.

$$\langle n(t) \rangle = \frac{e^{-\eta t \Delta \nu_f} \int_{-\infty}^t [G_{fr}^{(2)}(t') - G_{or}^{(2)}(t')] e^{\eta t' \Delta \nu_f} dt'}{\int_{-\infty}^{\infty} G_{fr}^{(2)}(t') dt'} \quad (6)$$

We estimated  $\eta$ , that includes the cavity losses per round trip and transmission through  $M_2$ , to be  $0.002 \pm 0.001$ .

When the cavity is exposed to the idler mode, a heralded single photon with decaying exponential envelope interacts with the cavity: the mean photon number in the cavity reaches a maximum of  $0.44 \pm 0.01$ . On the other hand, when the cavity is aligned in the signal mode we have a heralded single photon with an increasing exponential envelope interacting with the cavity; in this case,  $\langle n(t) \rangle$  reaches a maximum of  $0.76 \pm 0.01$ . As expected, the photon with the rising exponential waveform interacts more efficiently with the cavity. Following the analogy in [28], we expect this result to be extended to the

probability of absorption of a single photon by a single atom. In the case of interaction with a single atom, it will be necessary to also match the bandwidth of the transition. We have already demonstrated how it is possible to control the bandwidth of the photon generated by the cascade process by adjusting the optical density of the atomic medium [27].

In summary, we have demonstrated a method to transform a heralded single photon with a decaying exponential temporal envelope to a rising exponential envelope using a cavity. Using this method, we obtain single photons that resembles the time-reversed versions of photons from spontaneous decay process resonant to the D2 line of  $^{87}\text{Rb}$  atoms. Single photon states of this envelope and bandwidth would be useful for transferring information from photons back into atoms. [This time reversal technique can be used with photon pairs from other sources with time-ordered emission such as molecules and quantum dots](#), completing the toolbox necessary to interconnect [atoms stationary qubits](#) in a complex quantum information processing scenario.

We acknowledge the support of this work by the National Research Foundation & Ministry of Education in Singapore. [We thank Prof. Gerd Leuchs for useful discussions.](#)

---

\* Center for Quantum Technologies, 3 Science Drive 2, Singapore 117543

† Center for Quantum Technologies and Department of Physics, National University of Singapore, 3 Science Drive 2, Singapore 117543; christian.kurtsiefer@gmail.com

- [1] J. I. Cirac, P. Zoller, H. J. Kimble, and H. Mabuchi, *Phys. Rev. Lett.* **78**, 3221 (1997).
- [2] H. J. Kimble, *Nature* **453**, 1023 (2008).
- [3] L.-M. Duan, M. D. Lukin, J. I. Cirac, and P. Zoller, *Nature* **414**, 413 (2001).
- [4] A. Reiserer, N. Kalb, G. Rempe, and S. Ritter, *Nature* **508**, 237 (2014).
- [5] M. Stobińska, G. Alber, and G. Leuchs, *Europhys. Lett.* **86**, 14007 (2009).
- [6] Y. Wang, J. Minář, L. Sheridan, and V. Scarani, *Phys. Rev. A* **83**, 063842 (2011).
- [7] S. Ritter, C. Nolleke, C. Hahn, A. Reiserer, A. Neuzner, M. Uphoff, M. Mücke, E. Figueroa, J. Bochmann, and G. Rempe, *Nature* **484**, 195 (2012).
- [8] A. I. Lvovsky, B. Sanders, and W. Tittel, *Nat. Photonics* **3**, 706 (2009).
- [9] S. Zhang, C. Liu, S. Zhou, C.-S. Chu, M. M. T. Loy, and S. Du, *Phys. Rev. Lett.* **109**, 263601 (2012).
- [10] S. A. Aljunid, G. Maslennikov, Y. Wang, D. H. Lan, V. Scarani, and C. Kurtsiefer, *Phys. Rev. Lett.* **111**, 103001 (2013).
- [11] T. Volz, A. Reinhard, M. Winger, A. Badolato, K. J. Hennessy, E. L. Hu, and A. Imamoglu, *Nat Photon* **6**, 605 (2012).
- [12] R. Johne and A. Fiore, *Phys. Rev. A* **84**, 053850 (2011).
- [13] Y. L. A. Rezus, S. G. Walt, R. Lettow, A. Renn, G. Zumofen, S. Götzinger, and V. Sandoghdar, *Phys. Rev. Lett.* **108**, 093601 (2012).
- [14] J. Wenner, Y. Yin, Y. Chen, R. Barends, B. Chiaro, E. Jeffrey, J. Kelly, A. Megrant, J. Y. Mutus, C. Neill, et al., *Phys. Rev. Lett.* **112**, 210501 (2014).
- [15] P. Kolchin, C. Belthangady, S. Du, G. Y. Yin, and S. E. Harris, *Phys. Rev. Lett.* **101**, 103601 (2008).
- [16] G. K. Gulati, B. Srivathsan, B. Chng, A. Cerè, D. Matsukevich, and C. Kurtsiefer, [arXiv:1402.5800 \[PRA \(to be published\)\]](#).
- [17] M. Bader, S. Heugel, A. L. Chekhov, M. Sondermann, and G. Leuchs, *New Journal of Physics* **15**, 123008 (2013).
- [18] J. D. Franson, *Phys. Rev. A* **45**, 3126 (1992).
- [19] C. Santori, D. Fattal, M. Pelton, G. S. Solomon, and Y. Yamamoto, *Phys. Rev. B* **66**, 045308 (2002).
- [20] A. Dousse, J. Suffczynski, A. Beveratos, O. Krebs, A. Lemaître, I. Sagnes, J. Bloch, P. Voisin, and P. Senellart, *Nature* **466**, 217 (2010).
- [21] N. Akopian, N. H. Lindner, E. Poem, Y. Berlatzky, J. Avron, D. Gershoni, B. D. Gerardot, and P. M. Petroff, *Phys. Rev. Lett.* **96**, 130501 (2006).
- [22] D. V. Strekalov, A. V. Sergienko, D. N. Klyshko, and Y. H. Shih, *Phys. Rev. Lett.* **74**, 3600 (1995).
- [23] G. B. Lemos, V. Borish, G. D. Cole, S. Ramelow, R. Lapkiewicz, and A. Zeilinger, *Nature* **512**, 409 (2014).
- [24] H. H. Jen, *J. Phys. B: At. Mol. Opt. Phys.* **45**, 165504 (2012).
- [25] N. Hodgson and H. Weber, *Optical Resonators: Fundamentals, Advanced Concepts, and Applications* (Springer Verlag, 1997), ISBN 9783540761372.
- [26] G. S. Agarwal and S. D. Gupta, *Phys. Rev. A* **49**, 3954 (1994).
- [27] B. Srivathsan, G. K. Gulati, B. Chng, G. Maslennikov, D. Matsukevich, and C. Kurtsiefer, *Phys. Rev. Lett.* **111**, 123602 (2013).
- [28] S. Heugel, A. Villar, M. Sondermann, U. Peschel, and G. Leuchs, *Laser Physics* **20**, 100 (2010).